

A-12-00

PTO/SB/05 (12/97)

Please type a plus sign (+) inside this box → ☐Approved for use through 09/30/00. OMB 0651-0032
Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

UTILITY
PATENT APPLICATION
TRANSMITTAL

Attorney Docket No. SHAL 3.0-031 Total Pages 58

First Named Inventor or Application Identifier

Andrew V. Schally, et al.

Only for new nonprovisional applications under 37 CFR 1.53(b)

Express Mail Label No. EL 325030620US

APPLICATION ELEMENTS

1. MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO:

Assistant Commissioner for Patents
Box Patent Application
Washington, DC 202311. ☒ Fee Transmittal Form
(Submit an original, and a duplicate for fee processing)6. ☐ Microfiche Computer Program (Appendix)2. ☒ Specification [Total Pages 42]7. Nucleotide and/or Amino Acid Sequence Submission
(if applicable, all necessary)

- Descriptive title of the invention
- Cross References to Related Applications
- Statement Regarding Fed sponsored R & D
- Reference to Microfiche Appendix
- Background of the invention
- Brief Summary of the invention
- Brief Description of the Drawings (if filed)
- Detailed Description
- Claim(s)
- Abstract of the Disclosure

a. ☐ Computer Readable Copyb. ☐ Paper Copy (identical to computer copy)c. ☐ Statement verifying identity of above copies3. ☒ Drawing(s) (35 USC 113) [Total Sheets 5]
(executed copy)

4. Oath or Declaration [Total Pages 3]

a. ☐ Newly executed (original or copy)b. ☐ Copy from a prior application (37 CFR 1.63(d))
(for continuation/divisional with Box 17 completed)
[Note Box 5 below]i. ☐ DELETION OF INVENTOR(S)Signed statement attached deleting
inventor(s) named in the prior application,
see 37 CFR 1.63(d)(2) and 1.33(b).5. ☐ Incorporation By Reference (useable if Box 4b is checked)
The entire disclosure of the prior application, from which a
copy of the oath or declaration is supplied under Box 4b,
is considered as being part of the disclosure of the
accompanying application and is hereby incorporated by
reference therein.

ACCOMPANYING APPLICATION PARTS

8. ☒ Assignment Papers (cover sheet & document(s))9. ☐ 37 CFR 3.73(b) Statement ☐ Power of Attorney
(when there is an assignee)10. ☐ English Translation Document (if applicable)11. ☐ Information Disclosure
Statement (IDS)/PTO-1449 ☐ Copies of IDS
Citations12. ☒ Preliminary Amendment13. ☒ Return Receipt Postcard (MPEP 503)
(Should be specifically itemized)14. ☐ Small Entity ☐ Statement filed in prior application,
Statement(s) Status still proper and desired15. ☐ Certified Copy of Priority Document(s)
(if foreign priority is claimed)16. ☐ Other:

17. If a CONTINUING APPLICATION, check appropriate box and supply the requisite information:

☐ Continuation ☒ Divisional ☐ Continuation-in-part (CIP) of prior application No. 09 / 199,381

18. CORRESPONDENCE ADDRESS

☐ Customer Number or Bar Code Labelor ☒ Correspondence address below

(Insert Customer No. or Attach bar code label here)

NAME Omri M. Behr, Esq.

ADDRESS 325 Pierson Avenue

CITY Edison

STATE NJ

ZIP CODE 08837

COUNTRY USA

TELEPHONE 732-494-5240

FAX 732-494-0428

Barren Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Box Patent Application, Washington, DC 20231

04/11/00

JCS6 U.S. PTO

JCS6 U.S. PTO
09/547415
04/11/00

PATENT APPLICATION TRANSMITTAL LETTER

ATTORNEY'S DOCKET NO.

SHAL 3.0-031

TO THE COMMISSIONER OF PATENTS AND TRADEMARKS:

Transmitted herewith for filing is the patent application of Andrew V. Schally, et al.

for ANTAGONISTIC ANALOGS OF GH-RH INHIBITING IGF-I AND II

Enclosed are:

- ☒ 5 sheets sheets of drawing. ☐ Claim for small entity ☐ Independent
☐ an assignment of the invention to ☐ Small Business
☐ The Administrators of the Tulane Educational Fund ☐ Non-Profit

- ☐ a certified copy of a application.
☐ associate power of attorney.

CLAIMS AS FILED

FOR	NUMBER FILED	NUMBER EXTRA	RATE	FEE
TOTAL CLAIMS	3 -20 =	--	X \$ 9 \$ 18	--
INDEPENDENT CLAIMS	3 -3 =	--	X \$ 39 \$ 78	--
BASIC FEE 690.00/\$345.00				690.00
TOTAL FILING FEE				690.00

- ☐ Please charge my Deposit Account No. in the amount of \$
A duplicate copy of this sheet is enclosed.

- ☒ The Commissioner is hereby authorized to charge any additional fees which may be required at any time during the prosecution of this application without specific authorization, or credit any overpayment to Deposit Account No. 02-1750. A duplicate copy of this sheet is enclosed.

- ☒ A check in the amount of \$ 690.00 to cover the filing fee is enclosed.

date

Attorney of Record

Omri H. Behr

Regis. No. 22,940

Telephone (732) 494-5240

EXPRESS MAIL DEPOSIT
 RECEIVED
 DATE OF DEPOSIT
 THE SIGNATURE OF THE ATTORNEY OF RECORD
 IS BEING DEPOSITED WITH THE UNITED STATES
 POSTAL SERVICE "Express Mail First Office
 to Addressed" service under 37 CFR 1.101
 in the case indicated above and is
 addressed to the Commissioner of Patents
 and Trademarks, Washington, DC 20230

Omri H. Behr
 Reg. No. 22,940

SHAL 3.0-031

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Andrew V. Schally

Serial No.: Unallocated

Group Art Unit:

Filed: Herewith

Examiner:

For: ANTAGONISTIC ANALOGS OF GH-RH INHIBITING IGF-I AND II

Edison, NJ 08837

April 11, 2000

PRELIMINARY AMENDMENT

Hon. Commissioner of Patents and Trademarks

Washington, D. C. 20231

Sir:

Please amend the above-identified application as follows:

In the Claims

Cancel Claims 1 through 8.

Amend Claims 9 through 11 reading as follows:

9. (Amended) A [a]method of suppressing excessive levels of GH in a patient in need of same which comprises administering to said patient an effective amount of [a compound of Claim 1] a peptide selected from the group having the formulae:
X-R¹-R²-Asp-Ala-R⁵-R⁶-Thr-R⁸-R⁹-R¹⁰-Arg-R¹²-R¹³-R¹⁴-R¹⁵-R¹⁶-Leu-R¹⁸-R¹⁹-Arg-R²¹-R²²-
Leu-Gln-Asp-Ile-R²⁷-R²⁸-R²⁹-NH₂

wherein X is PhAc, IndAc, Ibu, Nac, 1- or 2-Npr, or Fpr,

R¹ is Tyr or His,
R² is D-Arg or D-Cit,
R⁵ is Ile or Val,
R⁶ is Phe, Nal or Phe(Y), in which Y=F, Cl, Br,
R⁸ is Asn, Gln, Ser, Thr, Ala, D-Asn, D-Gln, D-Ser, D-Thr, Abu, D-Abu, or Aib,
R⁹ is Arg, Har, Lys, Orn, D-Arg, D-Har, D-Lys, D-Orn, Cit, Nle, Tyr (Me), Ser, Ala
or Aib,
R¹⁰ is Tyr or Phe(Y), in which Y=H, F, Cl, Br, or OCH₃,
R¹² is Lys, D-Lys, or Orn,
R¹³ is Val or Nle,
R¹⁴ is Leu or Nle,
R¹⁵ is Gly, Ala, Abu, Nle or Gln,
R¹⁶ is Gln or Arg,
R¹⁸ is Ser or Nle,
R¹⁹ is Ala or Abu,
R²¹ is Lys or Orn,
R²² is Leu, Ala or Aib,
R²⁷ is Met, Leu, Nle, Abu, or D-Arg,
R²⁸ is Arg, D-Arg, Ser, Asn, Asp, Ala or Abu,
R²⁹ is Arg, D-Arg, Har or D-Har,
and pharmaceutically acceptable salts thereof.

10. (Amended) A[a] method of treating a patient having a cancer carrying
receptors for IGF-I or -II which comprises administering to said patient an effective amount
of a [compound of claim 1] peptide selected from the group having the formulae:
X-R¹-R²-Asp-Ala-R⁵-R⁶-Thr-R⁸-R⁹-R¹⁰-Arg-R¹²-R¹³-R¹⁴-R¹⁵-R¹⁶-Leu-R¹⁸-R¹⁹-Arg-R²¹-R²²-
Leu-Gln-Asp-Ile-R²⁷-R²⁸-R²⁹-NH₂

wherein X is PhAc, IndAc, Ibu, Nac, 1- or 2-Npr, or Fpr,
R¹ is Tyr or His,
R² is D-Arg or D-Cit,
R⁵ is Ile or Val,

R⁶ is Phe, Nal or Phe(Y), in which Y=F, Cl, Br,
R⁸ is Asn, Gln, Ser, Thr, Ala, D-Asn, D-Gln, D-Ser, D-Thr, Abu, D-Abu, or Aib,
R⁹ is Arg, Har, Lys, Orn, D-Arg, D-Har, D-Lys, D-Orn, Cit, Nle, Tyr (Me), Ser, Ala
or Aib,
R¹⁰ is Tyr or Phe(Y), in which Y=H, F, Cl, Br, or OCH₃,
R¹² is Lys, D-Lys, or Orn,
R¹³ is Val or Nle,
R¹⁴ is Leu or Nle,
R¹⁵ is Gly, Ala, Abu, Nle or Gln,
R¹⁶ is Gln or Arg,
R¹⁸ is Ser or Nle,
R¹⁹ is Ala or Abu,
R²¹ is Lys or Orn,
R²² is Leu, Ala or Aib,
R²⁷ is Met, Leu, Nle, Abu, or D-Arg,
R²⁸ is Arg, D-Arg, Ser, Asn, Asp, Ala or Abu,
R²⁹ is Arg, D-Arg, Har or D-Har,
and pharmaceutically acceptable salts thereof

11. (Amended) A[a] a method for inhibiting IGF-II levels in tumors (cancers) and the expression of mRNA for IGF-II in the same tumors, which comprises administering to said patient an effective amount of [a compound of Claim 1] a peptide selected from the group having the formulae:

X-R¹-R²-Asp-Ala-R⁵-R⁶-Thr-R⁸-R⁹-R¹⁰-Arg-R¹²-R¹³-R¹⁴-R¹⁵-R¹⁶-Leu-R¹⁸-R¹⁹-Arg-R²¹-R²²-
Leu-Gln-Asp-Ile-R²⁷-R²⁸-R²⁹-NH₂

wherein X is PhAc, IndAc, Ibu, Nac, 1- or 2-Npr, or Fpr,

R¹ is Tyr or His

R² is D-Arg or D-Cit,

R⁵ is Ile or Val,

R⁶ is Phe, Nal or Phe(Y), in which Y=F, Cl, Br,

R⁸ is Asn, Gln, Ser, Thr, Ala, D-Asn, D-Gln, D-Ser, D-Thr, Abu, D-Abu, or Aib.

R⁹ is Arg, Har, Lys, Orn, D-Arg, D-Har, D-Lys, D-Orn, Cit, Nle, Tyr (Me), Ser, Ala or Aib.

R¹⁰ is Tyr or Phe(Y), in which Y=H, F, Cl, Br, or OCH₃.

R¹² is Lys, D-Lys, or Orn.

R¹³ is Val or Nle.

R¹⁴ is Leu or Nle.

R¹⁵ is Gly, Ala, Abu, Nle or Gln.

R¹⁶ is Gln or Arg.

R¹⁸ is Ser or Nle.

R¹⁹ is Ala or Abu.

R²¹ is Lys or Orn.

R²² is Leu, Ala or Aib.

R²⁷ is Met, Leu, Nle, Abu, or D-Arg.

R²⁸ is Arg, D-Arg, Ser, Asn, Asp, Ala or Abu

R²⁹ is Arg, D-Arg, Har or D-Har.

and pharmaceutically acceptable salts thereof.

Respectfully submitted,


Omar M. Beltrami

Regis. No. 22,940

Tele: 732-494-5240

Shal31c12.plm

ANTAGONISTIC ANALOGS OF GH-RH INHIBITING IGF-I AND -II

FIELD OF INVENTION

This invention was made in part with Government support from the
5 Medical Research Service of the Veterans Affairs Department. The
Government has certain rights in this application.

The present invention relates to novel synthetic peptides which inhibit
the release of growth hormone from the pituitary in mammals, and to
10 therapeutic compositions containing these novel peptides.

BACKGROUND OF THE INVENTION

Growth Hormone ("GH") is a peptide having 191 amino acids which
stimulates the production of numerous different growth factors, e.g. insulin-
15 like growth factor I (IGF-I) and so promotes growth of numerous tissues
(skeleton, connective tissue, muscle and viscera) and physiological activities
(raising nucleic acid and protein synthesis and lipolysis, but lowering urea
secretion).

20 Release of GH is under the control of releasing and inhibiting factors
secreted by the hypothalamus. The primary releasing factor is growth
hormone releasing hormone ("GH-RH"); human growth hormone-releasing
hormone ("hGH-RH") is a peptide having 44 amino acids. The novel
peptides of the present invention relate to analogues of hGH-RH having only
25 residues 1 through 29 ("hGH-RH(1-29)NH₂"), i.e., to analogues of the
peptide which has the amino acid sequence:

Tyr-Ala-Asp-Ala-Ile⁶-Phe-Thr-Asn-Ser-Tyr¹⁰-Arg-Lys-Val-Leu-Gly¹⁵-
Gln-Leu-Ser-Ala-Arg²⁰-Lys-Leu-Leu-Gln-Asp²⁵-Ile-Met-Ser-Arg²⁹-NH₂

30 GH has been implicated in several diseases. One disease in which GH
is involved is acromegaly, in which excessive levels of GH are present. The

abnormally enlarged facial and extremity bones of this disease can be treated by administering a GH-RH antagonist.

Further diseases involving GH are diabetic retinopathy and diabetic nephropathy. The damage to the retina and kidneys respectively in these diseases, believed to be due to GH, results in blindness or reduction in kidney function. This damage can be prevented or slowed by administration of an effective GH-RH antagonist.

However, the main applications of GH-RH antagonists would be in the field of cancer (A.V. Schally et al, in *Growth Hormone Secretagogues in Clinical Practice*, eds. Bercu, B.B. & Walker, R.F., Dekker, New York, pp. 145-162, 1998). IGF-I and -II are potent mitogens for various cancers. By suppressing GH secretion, GH-RH antagonists decreases the synthesis of IGF-I in the liver and other tissues. GH-RH antagonists also reduce the autocrine and paracrine production of IGF-I and/or IGF-II by various tumors. In several experimental cancers, treatment with antagonists of GH-RH produces a reduction in IGF-I and -II, concomitant to inhibition of tumor growth.

In an effort to intervene in these disease and other conditions, some investigators have attempted to control GH levels by using somatostatin, one inhibitor of GH release. However, somatostatin, if administered alone, does not suppress GH or IGF-I levels to a desired degree. If administered in combination with a GH-RH antagonist, somatostatin would improve suppression of IGF-I levels much better.

Scientists have investigated various modifications of GH-RH to elucidate the relationship of the structure of GH-RH to its activity in an effort to provide synthetic congeners with improved agonistic or antagonistic properties. Thus, it was early established that GH-RH fragment comprising

residues 1 to 29, or GH-RH(1-29), is the minimum sequence necessary for biological activity. This fragment retains 50% or more of the potency of native GH-RH.

- 5 The first described GH-RH antagonist, [Ac-Tyr¹,D-Arg²]hGH-RH(1-29)NH₂, which is generally termed as the "standard antagonist" in the literature, was found to prevent the activation of rat anterior pituitary adenylate cyclase by hGH-RH(1-29)NH₂. The same peptide blocked the action of GH-RH on its receptors in the pituitary and hypothalamus, and
10 inhibited the pulsatile growth hormone secretion.

- A considerable number of patents and articles in the open literature disclose analogs of GH-RH which either act as agonists of GH-RH (i.e. act to stimulate the release of GH) or as antagonists of GH-RH (i.e. act to inhibit
15 the release of GH). Most of these peptides are derived from the GH-RH(1-29) peptide sequence, with specific structural modifications which account for their enhanced agonistic or antagonistic properties.

- Thus, US Patent 4,659,693 discloses GH-RH antagonistic analogs
20 which contain certain N,N'-dialkyl-omega-guanidino alpha-amino acyl residues in position 2 of the GH-RH(1-29) sequence.

- Published application WO 91/16923 reviews earlier attempts to alter the secondary structure of hGH-RH by modifying its amino acid sequence.
25 These earlier attempts include: replacing Tyr¹, Ala², Asp³ or Asn⁸ with their D-isomers; replacing Asn⁸ with L- or D-Ser, D-Arg, Asn, Thr, Gln or D-Lys; replacing Ser⁹ with Ala to enhance amphiphilicity of the region; and replacing Gly¹⁵ with Ala or Aib. When R² in the analogs is D-Arg, and R⁸, R⁹, and R¹⁵ are substituted as indicated above, antagonistic activity is said to
30 result. These antagonistic peptides are said to be suitable for administration as pharmaceutical compositions to treat conditions associated with

excessive levels of GH, e.g., acromegaly.

The antagonistic activity of the hGH-RH analogue "[Ser⁹-Ψ(CH₂-NH)-Tyr¹⁰]hGH-RH(1-29)" of US Patent 5,084,555 was said to result from the
5 pseudopeptide bond (i.e., a peptide bond reduced to a [CH₂-NH] linkage) between the R⁹ and R¹⁰ residues. However, the antagonistic properties of [Ser⁹-Ψ(CH₂-NH)-Tyr¹⁰]hGH-RH(1-29) were said to be inferior to the standard antagonist, [N-Ac-Tyr¹, D-Arg²]GH-RH(1-29)-NH₂.

10 US Patent 5,550,212, and US Patent Application 08/642,472, assigned to the same assignee as the present application, disclose analogs of hGH-RH(1-29)NH₂ said to have enhanced antagonistic properties and prolonged duration of action. These properties are believed to result from replacement of various amino acids and acylation with aromatic or nonpolar
15 acids at the N-terminus of GH-RH(1-29)NH₂. It is noted that in the above US Patent and US Patent Application, R⁹ is always Ser, R¹⁶ is Gln or an amino acid forming a lactam bridge (i.e. Glu), R²⁸ is Ser, Asn, Asp, Ala or Abu, and R²⁹ is Agm, Arg-NH₂, Arg-OH, Cit-NH₂, Cit-OH, Har-NH₂, Har-OH, or an amino acid forming a lactam bridge (i.e. Lys or Orn).

20

SUMMARY OF THE INVENTION

There is provided a novel series of synthetic analogs of hGH-RH(1-29)NH₂. These analogs inhibit the activity of endogenous hGH-RH, and therefore prevent the release of growth hormone. The stronger inhibitory
25 potencies of the new analogs, as compared to previously described ones, results from replacement of various amino acids.

Specifically, the invention relates to peptides comprising the formulae:
X-R¹-R²-Asp-Ala-R⁵-R⁶-Thr-R⁸-R⁹-R¹⁰-Arg-R¹²-R¹³-R¹⁴-R¹⁵-R¹⁶-Leu-R¹⁸-R¹⁹-Arg-
30 R²¹-R²²-Leu-Gln-Asp-Ile-R²⁷-R²⁸-R²⁹-NH₂

wherein X is PhAc, IndAc, Ibu, Nac, 1- or 2-Npr, or Fpr,

- R¹ is Tyr or His,
R² is D-Arg or D-Cit,
R⁵ is Ile or Val,
R⁶ is Phe, Nal or Phe(Y), in which Y = F, Cl, Br,
5 R⁸ is Asn, Gln, Ser, Thr, Ala, D-Asn, D-Gln, D-Ser, D-Thr, Abu, D-
Abu, or Aib,
R⁹ is Arg, Har, Lys, Orn, D-Arg, D-Har, D-Lys, D-Orn, Cit, Nle, Tyr
(Me), Ser, Ala or Aib,
R¹⁰ is Tyr or Phe(Y), in which Y = H, F, Cl, Br, or OCH₃,
10 R¹² is Lys, D-Lys, or Orn,
R¹³ is Val or Nle,
R¹⁴ is Leu or Nle,
R¹⁵ is Gly, Ala, Abu, Nle or Gln,
R¹⁶ is Gln or Arg,
15 R¹⁸ is Ser or Nle,
R¹⁹ is Ala or Abu,
R²¹ is Lys or Orn,
R²² is Leu, Ala or Aib,
R²⁷ is Met, Leu, Nle, Abu, or D-Arg,
20 R²⁸ is Arg, D-Arg, Ser, Asn, Asp, Ala or Abu,
R²⁹ is Arg, D-Arg, Har or D-Har,
and pharmaceutically acceptable salts thereof.

- Among the preferred embodiment are peptides wherein X is PhAc,
25 IndAc or Nac, R¹ is Tyr or His, R² is D-Arg, R⁵ is Ile, R⁶ is Phe(pCl), R⁸ is Asn
or Abu, R⁹ is Arg or Har, Lys, Orn, D-Arg, D-Har, D-Lys, D-Orn, Cit, Nle, or
Tyr (Me), R¹⁰ is Tyr or Tyr(Me), R¹² is Lys, R¹³ is Val or Nle, R¹⁴ is Leu or
Nle, R¹⁵ is Abu, Ala, or Nle, R¹⁶ is Gln or Arg, R¹⁸ is Ser or Nle, R¹⁹ is Ala or
Abu, R²¹ is Lys, R²⁷ is Nle or D-Arg,, R²⁸ is D-Arg, Arg, or Ser, R²⁹ is D-Arg,
30 Har or D-Har.

It is noted that the amino acid residues from 30 through 44 of the native GH-RH molecule do not appear to be essential to activity; nor does their identity appear to be critical. Therefore, it appears that the addition of some or all of these further amino acid residues to the C-terminus of the hGH-RH(1-29)-NH₂ analogues of the present invention will not affect the efficacy of these analogues as GH-RH antagonists. If some or all of these amino acids were added to the C-terminus of the hGH-RH(1-29)-NH₂ analogues, the added amino acid residues could be the same as residues 30 through 44 in the native hGH-RH sequence or reasonable equivalents.

Synthetic Methods.

The synthetic peptides are synthesized by a suitable method such as by exclusive solid phase techniques, by partial solid-phase techniques, by fragment condensation or by classical solution phase synthesis.

When the analogues of this invention are synthesized by solid-phase method, the C-terminus residue (here, R²⁹) is appropriately linked (anchored) to an inert solid support (resin) while bearing protecting groups for its alpha amino group (and, where appropriate, for its side chain functional group). After completion of this step, the alpha amino protecting group is removed from the anchored amino acid residue and the next amino acid residue, R²⁸, is added having its alpha amino group (as well as any appropriate side chain functional group) suitably protected, and so forth. The N-terminus protecting groups are removed after each residue is added, but the side chain protecting groups are not yet removed. After all the desired amino acids have been linked in the proper sequence, the peptide is cleaved from the support and freed from all side chain protecting group(s) under conditions that are minimally destructive towards residues in the sequence. This is followed by a careful purification and scrupulous characterization of the synthetic product, so as to ensure that the desired structure is indeed the one obtained.

It is particularly preferred to protect the alpha amino function of the amino acids during the coupling step with an acid or base sensitive protecting group. Such protecting groups should have the properties of being stable in the conditions of peptide linkage formation, while being readily removable without destruction of the growing peptide chain and without racemization of any of the chiral centers contained therein. Suitable alpha amino protecting groups are Boc and Fmoc.

Medical Applications.

10 The hGH-RH antagonist peptides, or salts of these peptides, may be formulated in pharmaceutical dosage forms containing effective amounts thereof and administered to humans or animal for therapeutic or diagnostic purposes. The peptides may be used to suppress GH levels and to treat conditions associated with excessive levels of GH, e.g., diabetic retinopathy and nephropathy, and acromegaly. Also provided are methods for treating these diseases by administration of a composition of the invention to an individual needing such treatment. The main uses of GH-RH antagonists are however, in the field of cancer, for example human cancers of the breast, lung, colon, brain, pancreas, and prostate where the receptors for IGF-I or IGF-II are present.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure I is a plot of volume changes of MXT mouse mammary cancers during treatment with certain GH-RH antagonists against days of treatment.

Figure II is a plot of volume changes of MDA-MB-468 human breast cancers in nude mice during treatment with certain GH-RH antagonists against days of treatment.

Figure III is a plot of volume changes of HT-29 human colon cancers in nude mice during treatment with certain GH-RH antagonists against days of treatment.

Figure IV is a plot of volume changes of U87MG human glioblastomas in nude mice during treatment with a GH-RH antagonist against days of treatment.

Figure V is a plot of volume changes of PC-3 human prostate cancers in nude mice during treatment with certain GH-RH antagonists against days of treatment.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

A. Abbreviations

10 The nomenclature used to define the peptides is that specified by the IUPAC-IUB Commission on Biochemical Nomenclature wherein, in accordance with conventional representation, the amino group at the N-terminus appears to the left and the carboxyl group at the C-terminus appears to the right. The term "natural amino acid" as used herein means
15 one of the common, naturally occurring L-amino acids found in naturally occurring proteins: Gly, Ala, Val, Leu, Ile, Ser, Thr, Lys, Arg, Asp, Asn, Glu, Gln, Cys, Met Phe, Tyr, Pro, Trp and His. When the natural amino acid residue has isomeric forms, it is the L-form of the amino acid that is represented herein unless otherwise expressly indicated.

20

Non-coded amino acids, or amino acid analogues, are also incorporated into the GH-RH antagonists. ("Non-coded" amino acids are those amino acids which are not among the approximately 20 natural amino acids found in naturally occurring peptides.) Among the non-coded amino
25 acids or amino acid analogues which may be used in the hGH-RH antagonist peptides are the following: by Abu is meant alpha amino butyric acid, by Aib is meant alpha amino isobutyric acid, by Har is meant homoarginine, by Nal is meant 2-naphthyl-alanine, by Nle is meant norleucine, and by Orn is meant ornithine. When these non-coded amino acids, or amino acid
30 analogues, have isomeric forms, it is the L-form of the amino acid that is represented unless otherwise expressly indicated.

Abbreviations used herein are:

	Abu	α -aminobutyric acid
	Ac	acetyl
	AcOH	acetic acid
5	Ac ₂ O	acetic anhydride
	Aib	α -aminoisobutyric acid
	Boc	tert.butyloxycarbonyl
	Bom	benzyloxymethyl
	2BrZ	2-bromo-benzyloxycarbonyl
10	cHx	cyclohexyl
	Cit	citrulline (2-amino-5-ureidovaleric acid)
	2ClZ	2-chloro-benzyloxycarbonyl
	DCM	dichloromethane
	DIC	N,N'-diisopropylcarbodiimide
15	DIEA	diisopropylethylamine
	DMF	dimethylformamide
	Fmoc	fluorenylmethyloxycarbonyl
	Fpr	3-phenylpropionyl
	GH	growth hormone
20	GH-RH	GH releasing hormone
	Har	homoarginine
	HBTU	2-(1H-Benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
	hGH-RH	human GH-RH
25	HOBt	1-hydroxybenzotriazole
	HPLC	high performance liquid chromatography
	Ibu	isobutryl
	IndAc	indole-3-acetyl
	MBHA	para-methylbenzhydrylamine
30	MeOH	methanol
	MeCN	acetonitrile

Nac	1-naphthylacetyl
Nal	2-naphthylalanine
NMM	N-methylmorpholine
Npr	naphthylpropionyl
5 PAM	phenylacetamidomethyl
Phe(pCl)	para-chloro-phenylalanine
PhAc	phenylacetyl
rGH-RH	rat GH-RH
RP-HPLC	reversed phase HPLC
10 TFA	trifluoroacetic acid
Tos	para-toluenesulfonyl
Tyr(Me)	tyrosine methylether
Z	benzyloxycarbonyl

15 B. The GH-RH Analogs

The hGH-RH analogues of the present invention were designed to increase the affinities of the peptides to the receptor, to improve metabolic stability and to maximize the amphiphilic secondary structure of the molecules. Many of these analogues cause very effective and long lasting inhibition of the GH release stimulated by hGH-RH(1-29)NH₂ in vitro and in vivo.

The following embodiments are specially preferred as having remarkable bioactivity:

- 25 [PhAc⁰, D-Arg², Phe(pCl)⁶, Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 1
- [IndAc⁰, D-Arg², Phe(pCl)⁶, Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 2
- [PhAc⁰, D-Arg², Phe(pCl)⁶, Har⁹, Tyr(Me)¹⁰, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 3
- 30 [PhAc⁰, D-Arg², Phe(pCl)⁶, Har⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-

29)NH₂ Peptide 4

[Nac⁰, D-Arg², Phe(pCl)⁶, Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 5

[PhAc⁰, D-Arg², Phe(pCl)⁶, Arg⁹, Tyr(Me)¹⁰, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-

5 RH(1-29)NH₂ Peptide 6

[PhAc⁰, His¹, D-Arg², Phe(pCl)⁶, Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 7

[Nac⁰, His¹, D-Arg², Phe(pCl)⁶, Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 8

10 [PhAc⁰, D-Arg², Phe(pCl)⁶, Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁹]hGH-RH(1-29)NH₂ Peptide 9

[PhAc⁰, D-Arg², Phe(pCl)⁶, Abu¹⁵, Arg¹⁶, Nle²⁷, D-Arg²⁹]hGH-RH(1-29)NH₂ Peptide 10

[PhAc⁰, D-Arg², Phe(pCl)⁶, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂

15 Peptide 11

[PhAc⁰, D-Arg², Phe(pCl)⁶, Nle⁹, Abu¹⁵, Nle²⁷, D-Arg²⁹]hGH-RH(1-29)NH₂

Peptide 12

[PhAc⁰, D-Arg², Phe(pCl)⁶, Nle¹³, Nle¹⁴, Abu¹⁵, Nle²⁷, D-Arg²⁹]hGH-RH(1-29)NH₂ Peptide 13

20 [PhAc⁰, D-Arg², Phe(pCl)⁶, Nle¹⁵, Nle²⁷, D-Arg²⁹]hGH-RH(1-29)NH₂

Peptide 14

[PhAc⁰, D-Arg², Phe(pCl)⁶, Abu¹⁵, Nle¹⁸, Nle²⁷, D-Arg²⁹]hGH-RH(1-29)NH₂

Peptide 15

[PhAc⁰, D-Arg², Phe(pCl)⁶, Tyr(Me)¹⁰, Abu¹⁵, Nle²⁷, D-Arg²⁹]hGH-RH(1-

25 29)NH₂ Peptide 16

[PhAc⁰, D-Arg², Phe(pCl)⁶, Abu⁸, Tyr(Me)¹⁰, Abu¹⁵, Nle²⁷, D-Arg²⁹]hGH-RH(1-29)NH₂ Peptide 17

[PhAc⁰, D-Arg², Phe(pCl)⁶, D-Abu⁸, Tyr(Me)¹⁰, Abu¹⁵, Nle²⁷, D-Arg²⁹]hGH-RH(1-29)NH₂ Peptide 18

30 [PhAc⁰, D-Arg², Phe(pCl)⁶, Tyr(Me)¹⁰, Abu¹⁵, D-Arg²⁷, Arg²⁸, D-Arg²⁹]hGH-RH(1-29)NH₂ Peptide 19

[PhAc⁰, D-Arg², Phe(pCl)⁶, Tyr(Me)⁹, Abu¹⁵, D-Arg²⁷, Arg²⁸, D-Arg²⁹]hGH-RH(1-29)NH₂ Peptide 20

[PhAc⁰, D-Arg², Phe(pCl)⁶, Abu¹⁵, D-Arg²⁷, Arg²⁸, D-Arg²⁹]hGH-RH(1-29)NH₂ Peptide 21

5 [PhAc⁰, D-Arg², Phe(pCl)⁶, Abu⁸, Tyr(Me)¹⁰, Abu¹⁵, D-Arg²⁷, Arg²⁸, D-Arg²⁹]hGH-RH(1-29)NH₂ Peptide 22

[PhAc⁰, D-Arg², Phe(pCl)⁶, D-Abu⁸, Tyr(Me)¹⁰, Abu¹⁵, D-Arg²⁷, Arg²⁸, D-Arg²⁹]hGH-RH(1-29)NH₂ Peptide 23

[PhAc⁰, D-Arg², Phe(pCl)⁶, Lys⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 24

[PhAc⁰, D-Arg², Phe(pCl)⁶, Orn⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 25

[PhAc⁰, D-Arg², Phe(pCl)⁶, D-Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 26

15 [PhAc⁰, D-Arg², Phe(pCl)⁶, D-Har⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 27

[PhAc⁰, D-Arg², Phe(pCl)⁶, D-Lys⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 28

[PhAc⁰, D-Arg², Phe(pCl)⁶, D-Orn⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 29

[PhAc⁰, D-Arg², Phe(pCl)⁶, Cit⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 30

Six very preferred embodiments have the formulae:

25 PhAc⁰-Tyr¹-D-Arg²-Asp³-Ala⁴-Ile⁵-Phe(pCl)⁶-Thr⁷-Asn⁸-Arg⁹-Tyr¹⁰-Arg¹¹-Lys¹²-Val¹³-Leu¹⁴-Abu¹⁵-Gln¹⁶-Leu¹⁷-Ser¹⁸-Ala¹⁹-Arg²⁰-Lys²¹-Leu²²-Leu²³-Gln²⁴-Asp²⁵-Ile²⁶-Nle²⁷-D-Arg²⁸-Har²⁹-NH₂ Peptide 1

IndAc⁰-Tyr¹-D-Arg²-Asp³-Ala⁴-Ile⁵-Phe(pCl)⁶-Thr⁷-Asn⁸-Arg⁹-Tyr¹⁰-Arg¹¹-Lys¹²-Val¹³-Leu¹⁴-Abu¹⁵-Gln¹⁶-Leu¹⁷-Ser¹⁸-Ala¹⁹-Arg²⁰-Lys²¹-Leu²²-Leu²³-Gln²⁴-Asp²⁵-

30 Ile²⁶-Nle²⁷-D-Arg²⁸-Har²⁹-NH₂ Peptide 2

PhAc⁰-Tyr¹-D-Arg²-Asp³-Ala⁴-Ile⁵-Phe(pCl)⁶-Thr⁷-Asn⁸-Har⁹-Tyr(Me)¹⁰-Arg¹¹-

- Lys¹²-Val¹³-Leu¹⁴-Abu¹⁵-Gln¹⁶-Leu¹⁷-Ser¹⁸-Ala¹⁹-Arg²⁰-Lys²¹-Leu²²-Leu²³-Gln²⁴-
 Asp²⁵-Ile²⁶-Nle²⁷-D-Arg²⁸-Har²⁹-NH₂ Peptide 3
- PhAc⁰-Tyr¹-D-Arg²-Asp³-Ala⁴-Ile⁵-Phe(pCl)⁶-Thr⁷-Asn⁸-Arg⁹-Tyr(Me)¹⁰-Arg¹¹-
 Lys¹²-Val¹³-Leu¹⁴-Abu¹⁵-Gln¹⁶-Leu¹⁷-Ser¹⁸-Ala¹⁹-Arg²⁰-Lys²¹-Leu²²-Leu²³-Gln²⁴-
 5 Asp²⁵-Ile²⁶-Nle²⁷-D-Arg²⁸-Har²⁹-NH₂ Peptide 6
- PhAc⁰-His¹-D-Arg²-Asp³-Ala⁴-Ile⁵-Phe(pCl)⁶-Thr⁷-Asn⁸-Arg⁹-Tyr¹⁰-Arg¹¹-Lys¹²-
 Val¹³-Leu¹⁴-Abu¹⁵-Gln¹⁶-Leu¹⁷-Ser¹⁸-Ala¹⁹-Arg²⁰-Lys²¹-Leu²²-Leu²³-Gln²⁴-Asp²⁵-
 Ile²⁶-Nle²⁷-D-Arg²⁸-Har²⁹-NH₂ Peptide 7
- Nac⁰-His¹-D-Arg²-Asp³-Ala⁴-Ile⁵-Phe(pCl)⁶-Thr⁷-Asn⁸-Arg⁹-Tyr¹⁰-Arg¹¹-Lys¹²-
 10 Val¹³-Leu¹⁴-Abu¹⁵-Gln¹⁶-Leu¹⁷-Ser¹⁸-Ala¹⁹-Arg²⁰-Lys²¹-Leu²²-Leu²³-Gln²⁴-Asp²⁵-
 Ile²⁶-Nle²⁷-D-Arg²⁸-Har²⁹-NH₂ Peptide 8

Under well-established convention, these may be abbreviated as follows:

- 15 [PhAc⁰, D-Arg², Phe(pCl)⁶, Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-
 29)NH₂ Peptide 1
- [IndAc⁰, D-Arg², Phe(pCl)⁶, Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-
 29)NH₂ Peptide 2
- [PhAc⁰, D-Arg², Phe(pCl)⁶, Har⁹, Tyr(Me)¹⁰, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-
 20 RH(1-29)NH₂ Peptide 3
- [PhAc⁰, D-Arg², Phe(pCl)⁶, Arg⁹, Tyr(Me)¹⁰, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-
 RH(1-29)NH₂ Peptide 6
- [PhAc⁰, His¹, D-Arg², Phe(pCl)⁶, Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-
 29)NH₂ Peptide 7
- 25 [Nac⁰, His¹, D-Arg², Phe(pCl)⁶, Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-
 29)NH₂ Peptide 8

C. Method of Preparation

1. Overview of Synthesis

- 30 The peptides are synthesized by suitable methods such as by
 exclusive solid phase techniques, by partial solid-phase techniques, by

fragment condensation or by classical solution phase synthesis. For example, the techniques of exclusive solid-phase synthesis are set forth in the textbook "Solid Phase Peptide Synthesis", J.M. Stewart and J.D. Young, Pierce Chem. Company, Rockford, 111, 1984 (2nd. ed.), and M. Bodanszky, "Principles of Peptide Synthesis", Springer Verlag, 1984. The hGH-RH antagonist peptides are preferably prepared using solid phase synthesis, such as that generally described by Merrifield, J. Am. Chem. Soc., 85 p. 2149 (1963), although other equivalent chemical syntheses known in the art can also be used as previously mentioned.

10

The synthesis is carried out with amino acids that are protected at their alpha amino group. Urethane type protecting groups (Boc or Fmoc) are preferably used for the protection of the alpha amino group. The preferred protecting group is Boc.

15

In solid phase synthesis, the N-alpha-protected amino acid moiety which forms the aminoacyl group of the final peptide at the C-terminus is attached to a polymeric resin support via a chemical link. After completion of the coupling reaction, the alpha amino protecting group is selectively removed to allow subsequent coupling reactions to take place at the amino-terminus, preferably with 50% TFA in DCM when the N-alpha-protecting group is Boc. The remaining amino acids with similarly Boc-protected alpha amino groups are coupled stepwise to the free amino group of the preceding amino acid on the resin to obtain the desired peptide sequence. Because the amino acid residues are coupled to the alpha amino group of the C-terminus residue, growth of the synthetic hGH-RH analogue peptides begins at the C terminus and progresses toward the N-terminus. When the desired sequence has been obtained, the peptide is acylated at the N-terminus, and it is removed from the support polymer.

30

Each protected amino acid is used in excess (2.5 or 3 equivalents) and the coupling reactions are usually carried out in DCM, DMF or mixtures thereof. The extent of completion of the coupling reaction is monitored at each stage by the ninhydrin reaction. In cases where incomplete coupling is determined, the coupling procedure is repeated, or a capping by acetylation of unreacted amino groups is carried out, before removal of the alpha amino protecting group prior to the coupling of the next amino acid.

A typical synthesis cycle is shown in Table I.

TABLE I		
Protocol for a Typical Synthetic Cycle Using Boc-strategy		
Step	Reagent	Mixing Time (min)
1. Deprotection	50% TFA in DCM	5 + 25
	DCM wash	1
	2-propanol wash	1
2. Neutralization	5% DIEA in DCM	1
	DCM wash	1
	MeOH wash	1
	5% DIEA in DCM	3
	MeOH wash	1
	DCM wash (3 times)	1-1
3. Coupling	3 equiv. Boc-amino acid in DCM or DMF + 3 equiv. DIC or the preformed HOBt ester of the Boc-amino acid	60
	MeOH wash	2
	DCM wash	2
4. Acetylation (if appropriate)	Ac ₂ O in DCM (30%)	10 + 20
	MeOH wash (3 times)	2
	DCM wash (3 times)	2

After completion of the synthesis, the cleavage of the peptide from the resin can be effected using procedures well known in peptide chemistry.

Some of the amino acid residues of the peptides have side chain functional groups which are reactive with reagents used in coupling or

deprotection. When such side chain groups are present, suitable protecting groups are joined to these functional groups to prevent undesirable chemical reactions occurring during the reactions used to form the peptides. The following general rules are followed in selecting a particular side chain

5 protecting group: (a) the protecting group preferably retains its protecting properties and is not split off under coupling conditions, (b) the protecting group should be stable under conditions for removing the alpha amino protecting group at each step of the synthesis, (c) the side chain protecting group must be removable upon the completion of the synthesis of the

10 desired amino acid sequence, under reaction conditions that will not undesirably alter the peptide chain.

The reactive side chain functional groups are preferably protected as follows: benzyl for Thr and Ser; 2-bromo-benzyloxycarbonyl for Tyr; p-

15 toluene-sulfonyl or nitro for Arg and Har; 2-chlorobenzyloxycarbonyl or fluorenylmethyloxycarbonyl for Lys, Orn; benzyloxymethyl for His; and cyclohexyl or fluorenylmethyl for Asp and Glu. The side chains of Asn and Gln are unprotected.

20 3. Stepwise Coupling of Amino Acid Residues to the Support Polymer

The hGH-RH antagonist peptides may be synthesized on a variety of support polymers, i.e. MBHA, Merrifield, PAM or Wang resins. When N-alpha-Boc protected amino acids are used for synthesis, the preferred resin

25 is MBHA. In this case, peptides with an amidated C-terminus are obtained upon cleavage from the support phase.

First, the C-terminal amino acid is attached to the neutralized MBHA resin, and then the subsequent amino acid couplings are carried out. Each

30 protected amino acid is coupled in about a three-fold molar excess, with respect to resin-bound free amino residues, and the coupling may be carried

out in a medium such as DMF: CH_2Cl_2 (1:1) or in DMF or CH_2Cl_2 alone. The selection of an appropriate coupling reagent is within the skill of the art. Particularly suitable as coupling reagents are N,N'-diisopropyl carbodiimide (DIC), or HBTU combined with HOBt. The success of the coupling reaction at each stage of the synthesis is preferably monitored by the ninhydrin reaction. In cases where incomplete coupling occurs, either the coupling procedure is repeated, or the resin-bound unreacted amino residues are acetylated using $\text{Ac}_2\text{O}/\text{DCM}$, before removal of the alpha amino protecting group.

10

Final acylation of the N-terminus of the peptide is done in the same way as the previous couplings, with the difference that the appropriate carboxylic acid is used instead of an amino acid.

15

4. Removal of the Peptide from the Support Polymer.

When the synthesis is complete, the peptide is cleaved from the support phase. Removal of the peptide from the resin is performed by treatment with a reagent such as liquid hydrogen fluoride which also cleaves all remaining side chain protecting groups.

20

Suitably, the dried and protected peptide-resin is treated with a mixture consisting of 1.0 mL m-cresol and 10 mL anhydrous hydrogen fluoride per gram of peptide-resin for 60 min at 0°C to cleave the peptide from the resin as well as to remove all side chain protecting groups. After the removal of the hydrogen fluoride under a stream of nitrogen and vacuum, the free peptides are precipitated with ether, filtered, washed with ether and ethyl acetate, extracted with 50% acetic acid, and lyophilized.

30

5. Purification

The purification of the crude peptides can be effected using procedures well known in peptide chemistry. For example, purification may

be performed on a MacRabbit HPLC system (Rainin Instrument Co. Inc., Woburn, MA) with a Knauer UV Photometer and a Kipp and Zonen BD40 Recorder using a Vydac 218TP5010 reversed-phase column (10 x 250 mm, packed with C18 silica gel, 300 Å pore size, 5 µm particle size) (The
5 Separations Group Inc., Hesperia, CA). The column is eluted with a solvent system consisting of (A) 0.1% aqueous TFA and (B) 0.1% TFA in 70% aqueous MeCN in a linear gradient mode (e.g., 30-55% B in 120 min). The eluent is monitored at 220 nm, and fractions are examined by analytical HPLC using a Hewlett-Packard Model HP-1090 liquid chromatograph and
10 pooled to give maximum purity. Analytical HPLC is carried out on a Vydac 218TP52 reversed-phase column (2 x 250 mm, C18, 300 Å, 5 µm) using isocratic elution with a solvent system consisting of (A) and (B) defined above. The peaks are monitored at 220 and 280 nm. The peptides are judged to be substantially (>95%) pure by analytical HPLC. The expected
15 amino acid composition is also confirmed by amino acid analysis.

D. Pharmaceutical Composition

The peptides of the invention may be administered in the form of pharmaceutically acceptable, nontoxic salts, such as acid addition salts.
20 Illustrative of such acid addition salts are hydrochloride, hydrobromide, sulphate, phosphate, fumarate, gluconate, tannate, maleate, acetate, citrate, benzoate, succinate, alginate, pamoate, malate, ascorbate, tartarate, and the like. Particularly preferred antagonists are salts of low solubility, e.g., pamoate salts and the like. These exhibit long duration of activity.

25

The compounds of the present invention are suitably administered to subject humans or animals s.c., i.m., or i.v; intranasally or by pulmonary inhalation; or in a depot form (e.g., microcapsules, microgranules, or cylindrical rod like implants) formulated from a biodegradable suitable
30 polymer (such as D,L-lactide-coglycolide), the former two depot modes being preferred. Other equivalent modes of administration are also within

the scope of this invention, i.e., continuous drip, depot injections, infusion pump and time release modes such as microcapsules and the like. Administration is in any physiologically acceptable injectable carrier, physiological saline being acceptable, though other carriers known to the art
5 may also be used.

The peptides are preferably administered parenterally, intramuscularly, subcutaneously or intravenously with a pharmaceutically acceptable carrier such as isotonic saline. Alternatively, the peptides may be administered as
10 an intranasal spray with an appropriate carrier or by pulmonary inhalation. One suitable route of administration is a depot form formulated from a biodegradable suitable polymer, e.g., poly-D,L-lactide-coglycolide as microcapsules, microgranules or cylindrical implants containing dispersed antagonistic compounds.

15

The amount of peptide needed depends on the mode of administration and the intended result. In general, the dosage range is between 1-100 $\mu\text{g/kg}$ of body weight of the host per day.

20 E. Therapeutic Uses of GH-RH Antagonists

hGH-RH antagonists can be used in treatment of conditions caused by excess growth hormone, for example acromegaly, which is manifested by an abnormal enlargement of the bones of the face and extremities. The GH-RH antagonists may also be used to treat diabetic nephropathy (the main
25 cause of blindness in diabetics) and diabetic retinopathy, in which damage to the eye and kidney respectively is thought to be due to GH.

The hGH-RH antagonists are designed to block the binding and therefore the action of GH-RH, which stimulates the secretion of GH, which
30 in turn stimulates production of IGF-I. GH-RH antagonists may be administered alone or together with somatostatin analogues, a combination

which more completely suppresses IGF-I levels. It is advantageous to administer antagonists of GH-RH rather than somatostatin due to the fact that GH-RH antagonists may be utilized in situations where target sites do not have somatostatin receptors.

5

However, the main applications of GH-RH antagonists are in the field of cancer. This is based on the following considerations: GH-RH antagonists are designed to block the binding and therefore the action of GH-RH, which stimulates the secretion of GH, which in turn stimulates production of insulin-like growth factor I (IGF-I) also called somatomedin-C. The involvement of IGF-I (somatomedin-C) in breast cancer, prostate cancer, colon cancer, bone tumors and other malignancies is well established, and somatostatin analogues alone do not adequately suppress GH and IGF-I levels. A complete suppression of IGF-I levels or secretion is required for a better inhibition of tumor growth. Autocrine production of IGF-I by various tumors could be also under control of GH-RH and might therefore be inhibited by GH-RH antagonists. GH-RH antagonists might also inhibit the production of IGF-I. A more detailed theoretical background of the applications of GH-RH in the field of oncology (cancer) is as follows: The receptors for IGF-I are present in primary human breast cancers, prostate cancers, lung cancers, colon cancers, brain tumors, pancreatic cancers, and in renal cell carcinomas.

The presence of IGF-I receptors in these tumors appears to be related to malignant transformation and proliferations of these cancers. IGF-I can act as endocrine, paracrine or autocrine growth factor for various human cancers, that is the growth of these neoplasms is dependent on IGF-I. GH-RH antagonists by suppressing GH secretion would lower the production of IGF-I. Since IGF-I stimulates growth of these various neoplasms (cancers), the lowering of circulating IGF-I levels should lead to tumor growth inhibition. It is possible that GH-RH antagonists could also lower paracrine

or autocrine production of IGF-I by the tumors, which should also lead to inhibition of cancer proliferation. These views are in accordance with modern concepts of clinical oncology. GH-RH antagonists should be given alone or together with somatostatin analogues and a combination would
5 achieve a more complete suppression of IGF-I levels, elimination of tissue IGF-I levels, e.g., in human osteosarcomas, as well as breast cancer, colon cancer, prostate cancer, and non-small cell lung cancer (non-SCLC).

The advantage of GH-RH antagonists over somatostatin analogues is
10 based on the fact that GH-RH antagonists may be utilized for suppression of tumors which do not have somatostatin receptors, for example human osteogenic sarcomas.

Antagonistic analogs of GH-RH have been shown to suppress growth
15 of various tumors *in vivo*. This effect is exerted in part through inhibition of the GHRH-GH-IGF-I axis. Nevertheless, autocrine/paracrine control of proliferation by IGF-II is also a major factor in many tumors. Interference with this autocrine growth-stimulating pathway offers an approach to tumor control. Antagonistic analogs of GH-RH, MZ-4-71 {[Ibu⁰, Tyr¹, D-Arg²,
20 Abu¹⁵, Nle²⁷]hGH-RH(1-28) Agm} and MZ-5-156 {[PhAc⁰, D-Arg², Abu¹⁵, Nle²⁷]hGH-RH(1-28) Agm} significantly inhibited the rate of proliferation of mammary (MDA-MB-468, ZR-75-1), prostatic (PC-3 and DU-145), and pancreatic (MiaPaCa-2, SW-1990 and Capan-2) cancer cell lines *in vitro* as shown by colorimetric and [³H]-thymidine incorporation tests, reduced the
25 expression of IGF-II mRNA in the cells and the concentration of IGF-II secreted into the culture medium. The same GH-RH antagonists produced similar results *in vivo* (inhibition of proliferation and reduction of IGF-II production) for prostate tumors (PC-3, DU-145), renal adenocarcinoma (Caki-I) and non-small cell lung carcinoma (H157). These findings suggest
30 that antagonistic analogs of GH-RH can inhibit tumor growth not only by inhibiting the GHRH-GH-IGF-I axis, but also by reducing the IGF-II production

in certain tumor cells, thus interrupting its autocrine regulatory pathway.

The present invention is described in connection with the following examples which are set forth for the purposes of illustration only. In the 5 examples, optically active protected amino acids in the L-configuration are used except where specifically noted.

The following Examples set forth suitable methods of synthesizing the novel GH-RH antagonists by the solid-phase technique.

10

EXAMPLE I

PhAc⁰-Tyr¹-D-Arg²-Asp³-Ala⁴-Ile⁵-Phe(pCl)⁶-Thr⁷-Asn⁸-Arg⁹-Tyr¹⁰-Arg¹¹-Lys¹²-
Val¹³-Leu¹⁴-Abu¹⁵-Gln¹⁶-Leu¹⁷-Ser¹⁸-Ala¹⁹-Arg²⁰-Lys²¹-Leu²²-Leu²³-Gln²⁴-Asp²⁵-
Ile²⁶-Nle²⁷-D-Arg²⁸-Har²⁹-NH₂ (Peptide 1)

5 { [PhAc⁰, D-Arg², Phe(pCl)⁶, Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ }

The synthesis is conducted in a stepwise manner using manual solid phase peptide synthesis equipment. Briefly, para-methylbenzhydrylamine (MBHA) resin (Bachem, California) (720 mg, 0.50 mmole) is neutralized with
10 5% DIEA in CH₂Cl₂ and washed according to the protocol described in Table I. The solution of Boc-Har(NO₂)-OH (500 mg, 1.5 mmole) in DMF-CH₂Cl₂ (1:1) is shaken with the neutralized resin and DIC (235 μL, 1.5 mmole) in a manual solid phase peptide synthesis apparatus for 1 hour. After the completion of the coupling reaction is proved by negative ninhydrin test,
15 deprotection with 50% TFA in CH₂Cl₂, and neutralization with 5% DIEA in CH₂Cl₂, the peptide chain is built stepwise by coupling the following protected amino acids in the indicated order on the resin to obtain the desired peptide sequence: Boc-D-Arg(Tos)-OH, Boc-Nle-OH, Boc-Ile-OH, Boc-Asp(OcHx)-OH, Boc-Gln-OH, Boc-Leu-OH, Boc-Leu-OH, Boc-Lys(2ClZ)-OH,
20 Boc-Arg(Tos)-OH, Boc-Ala-OH, Boc-Ser(Bzl)-OH, Boc-Leu-OH, Boc-Gln-OH, Boc-Abu-OH, Boc-Leu-OH, Boc-Val-OH, Boc-Lys(2ClZ)-OH, Boc-Arg(Tos)-OH, Boc-Tyr(2BrZ)-OH, Boc-Arg(Tos)-OH, Boc-Asn-OH, Boc-Thr(Bzl)-OH, Boc-Phe(pCl)-OH, Boc-Ile-OH, Boc-Ala-OH, Boc-Asp(OcHx)-OH, Boc-D-Arg(Tos)-OH, Boc-Tyr(2BrZ)-OH.

25

These protected amino acid residues (also commonly available from Bachem Co.) are represented above according to a well accepted convention. The suitable protecting group for the side chain functional group of particular amino acids appears in parentheses. The OH groups in the
30 above formulae indicate that each residue's carboxyl terminus is free.

The protected amino acids (1.5 mmole each) are coupled with DIC (235 μ L, 1.5 mmole) with the exceptions of Boc-Asn-OH and Boc-Gln-OH which are coupled with their preformed HOBt esters. After removal of the N^o-Boc protecting group from Tyr¹, the peptide is acylated with phenylacetic acid (PhAc) (272 mg, 2 mmole) using DIC (313 μ L, 2 mmole).

In order to cleave the peptide from the resin and deprotect it, the dried peptide resin (2.18 g) is stirred with 2 mL m-cresol and 20 mL hydrogen fluoride (HF) at 0°C for 1 hour. After evaporation of the HF under vacuum, the remnant is washed with dry diethyl ether and ethyl acetate. The cleaved and deprotected peptide is dissolved in 50 % acetic acid and separated from the resin by filtration. After dilution with water and lyophilization, 1.51 g crude product is obtained.

The crude peptide is checked by analytical HPLC using a Hewlett-Packard Model HP-1090 liquid chromatograph with a Vydac 218TP52 reversed-phase column (2 x 250 mm, packed with C18 silica gel, 300 Å pore size, 5 μ m particle size) (The Separations Group Inc., Hesperia, CA) and linear gradient elution, (e.g., 40-70% B) with a solvent system consisting of (A) 0.1% aqueous TFA and (B) 0.1% TFA in 70% aqueous MeCN. 500 mg of the crude peptide is dissolved in AcOH/H₂O, stirred, filtered and applied on a Beckman Ultraprep ODS column (21.2 x 150 mm, packed with C18 silica gel, 300 Å pore size, 10 μ m particle size). The column is eluted with a solvent system described above in a linear gradient mode (e.g., 30-55% B in 120 min); flow rate 6 mL/min. The eluent is monitored at 220 nm, and fractions are examined by analytical HPLC. Fractions with purity higher than 95% are pooled and lyophilized to give 98 mg pure product. The analytical HPLC is carried out on a Vydac C18 reversed-phase column described above using isocratic elution with a solvent system described above with a flow rate of 0.2 mL/min. The peaks are monitored at 220 and 280 nm. The product is judged to be substantially

(>95%) pure by analytical HPLC. Molecular mass is checked by electro-spray mass spectrometry, and the expected amino acid composition is confirmed by amino acid analysis.

5

EXAMPLE II

PhAc⁰-Tyr¹-D-Arg²-Asp³-Ala⁴-Ile⁵-Phe(pCl)⁶-Thr⁷-Asn⁸-Har⁹-Tyr(Me)¹⁰-Arg¹¹-Lys¹²-Val¹³-Leu¹⁴-Abu¹⁵-Gln¹⁶-Leu¹⁷-Ser¹⁸-Ala¹⁹-Arg²⁰-Lys²¹-Leu²²-Leu²³-Gln²⁴-Asp²⁵-Ile²⁶-Nle²⁷-D-Arg²⁸-Har²⁹-NH₂ (Peptide 3)

{[PhAc⁰, D-Arg², Phe(pCl)⁶, Har⁹, Tyr(Me)¹⁰, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂}

The synthesis is conducted in a stepwise manner using manual solid phase peptide synthesis equipment. Briefly, para-methylbenzhydrylamine (MBHA) resin (Bachem, California) (100 mg, 0.070 mmole) is neutralized with 5% DIEA in CH₂Cl₂ and washed according to the protocol described in Table I. The solution of Boc-Har(NO₂)-OH (83mg, 0.25 mmole) in DMF-CH₂Cl₂ (1:1) is shaken with the neutralized resin and DIC (44 μL, 0.275 mmole) in a manual solid phase peptide synthesis equipment for 1 hour. After the completion of the coupling reaction is proved by negative ninhydrin test, deprotection with 50% TFA in CH₂Cl₂, and neutralization with 5% DIEA in CH₂Cl₂, the peptide chain is built stepwise by coupling the following protected amino acids in the indicated order on the resin to obtain the desired peptide sequence: Boc-D-Arg(Tos)-OH, Boc-Nle-OH, Boc-Ile-OH, Boc-Asp(OcHx)-OH, Boc-Gln-OH, Boc-Leu-OH, Boc-Lys(2ClZ)-OH, Boc-Arg(Tos)-OH, Boc-Ala-OH, Boc-Ser(Bzl)-OH, Boc-Leu-OH, Boc-Gln-OH, Boc-Abu-OH, Boc-Leu-OH, Boc-Val-OH, Boc-Lys(2ClZ)-OH, Boc-Arg(Tos)-OH, Boc-Tyr(Me)-OH, Boc-Har(NO₂)-OH, Boc-Asn-OH, Boc-Thr(Bzl)-OH, Boc-Phe(pCl)-OH, Boc-Ile-OH, Boc-Ala-OH, Boc-Asp(OcHx)-OH, Boc-D-Arg(Tos)-OH, Boc-Tyr(2BrZ)-OH.

These protected amino acid residues (also commonly available from Bachem Co.) are represented above according to a well accepted con-

vention. The suitable protecting group for the side chain functional group of particular amino acids appears in parentheses. The OH groups in the above formulae indicate that each residue's carboxyl terminus is free.

5 The protected amino acids (0.25 mmole each) are coupled with DIC (44 μ L, 0.275 mmole) with the exceptions of Boc-Asn-OH and Boc-Gln-OH which are coupled with their preformed HOBt esters. After removal of the N^o-Boc protecting group from Tyr¹, the peptide is acylated with phenylacetic acid (PhAc) (54 mg, 0.4 mmole) using DIC (70 μ L, 0.44 mmole).

10

In order to cleave the peptide from the resin and deprotect it, the dried peptide resin (206 mg) is stirred with 0.5 mL m-cresol and 5 mL hydrogen fluoride (HF) at 0°C for 1 hour. After evaporation of the HF under vacuum, the remnant is washed with dry diethyl ether and ethyl acetate.

15 The cleaved and deprotected peptide is dissolved in 50 % acetic acid and separated from the resin by filtration. After dilution with water and lyophilization, 112 mg crude product is obtained.

The crude peptide is checked by analytical HPLC using a Hewlett-
20 Packard Model HP-1090 liquid chromatograph with a Vydac 218TP52 reversed-phase column (2 x 250 mm, packed with C18 silica gel, 300 Å pore size, 5 μ m particle size) (The Separations Group Inc., Hesperia, CA) and linear gradient elution, (e.g., 40-70% B) with a solvent system consisting of (A) 0.1% aqueous TFA and (B) 0.1% TFA in 70% aqueous
25 MeCN. 80 mg of the crude peptide is dissolved in AcOH/H₂O, stirred, filtered and applied on a Vydac 218TP5010 column (10 x 250 mm) packed with C8 silica gel. The column is eluted with a solvent system described above in a linear gradient mode (e.g., 30-55% B in 120 min); flow rate 2mL/min. The eluent is monitored at 220 nm, and fractions are examined
30 by analytical HPLC. Fractions with purity higher than 95% are pooled and lyophilized to give 9.6 mg pure product. The analytical HPLC is carried out

on a Vydac C18 reversed-phase column described above using isocratic elution with a solvent system described above with a flow rate of 0.2 mL/min. The peaks are monitored at 220 and 280 nm. The product is judged to be substantially (> 95%) pure by analytical HPLC. Molecular mass is checked by electrospray mass spectrometry, and the expected amino acid composition is confirmed by amino acid analysis.

Peptide 2, and Peptides 4 through 30 are synthesized in the same manner as Peptide 1 and Peptide 3, except that these peptides also contain other substitutions, to yield:

- [IndAc⁰, D-Arg², Phe(pCl)⁶, Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 2
- [PhAc⁰, D-Arg², Phe(pCl)⁶, Har⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 4
- 15 [Nac⁰, D-Arg², Phe(pCl)⁶, Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 5
- [PhAc⁰, D-Arg², Phe(pCl)⁶, Arg⁹, Tyr(Me)¹⁰, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 6
- [PhAc⁰, His¹, D-Arg², Phe(pCl)⁶, Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 7
- 20 [Nac⁰, His¹, D-Arg², Phe(pCl)⁶, Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 8
- [PhAc⁰, D-Arg², Phe(pCl)⁶, Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁹]hGH-RH(1-29)NH₂ Peptide 9
- 25 [PhAc⁰, D-Arg², Phe(pCl)⁶, Abu¹⁵, Arg¹⁶, Nle²⁷, D-Arg²⁹]hGH-RH(1-29)NH₂ Peptide 10
- [PhAc⁰, D-Arg², Phe(pCl)⁶, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 11
- [PhAc⁰, D-Arg², Phe(pCl)⁶, Nle⁹, Abu¹⁵, Nle²⁷, D-Arg²⁹]hGH-RH(1-29)NH₂ Peptide 12
- 30 [PhAc⁰, D-Arg², Phe(pCl)⁶, Nle¹³, Nle¹⁴, Abu¹⁵, Nle²⁷, D-Arg²⁹]hGH-RH(1-

29)NH₂ Peptide 13

[PhAc⁰, D-Arg², Phe(pCl)⁶, Nle¹⁵, Nle²⁷, D-Arg²⁹]hGH-RH(1-29)NH₂

Peptide 14

[PhAc⁰, D-Arg², Phe(pCl)⁶, Abu¹⁵, Nle¹⁸, Nle²⁷, D-Arg²⁹]hGH-RH(1-29)NH₂

5 Peptide 15

[PhAc⁰, D-Arg², Phe(pCl)⁶, Tyr(Me)¹⁰, Abu¹⁵, Nle²⁷, D-Arg²⁹]hGH-RH(1-29)NH₂ Peptide 16

[PhAc⁰, D-Arg², Phe(pCl)⁶, Abu⁸, Tyr(Me)¹⁰, Abu¹⁵, Nle²⁷, D-Arg²⁹]hGH-RH(1-29)NH₂ Peptide 17

10 [PhAc⁰, D-Arg², Phe(pCl)⁶, D-Abu⁸, Tyr(Me)¹⁰, Abu¹⁵, Nle²⁷, D-Arg²⁹]hGH-RH(1-29)NH₂ Peptide 18

[PhAc⁰, D-Arg², Phe(pCl)⁶, Tyr(Me)¹⁰, Abu¹⁵, D-Arg²⁷, Arg²⁸, D-Arg²⁹]hGH-RH(1-29)NH₂ Peptide 19

[PhAc⁰, D-Arg², Phe(pCl)⁶, Tyr(Me)⁹, Abu¹⁵, D-Arg²⁷, Arg²⁸, D-Arg²⁹]hGH-

15 RH(1-29)NH₂ Peptide 20

[PhAc⁰, D-Arg², Phe(pCl)⁶, Abu¹⁵, D-Arg²⁷, Arg²⁸, D-Arg²⁹]hGH-RH(1-29)NH₂ Peptide 21

[PhAc⁰, D-Arg², Phe(pCl)⁶, Abu⁸, Tyr(Me)¹⁰, Abu¹⁵, D-Arg²⁷, Arg²⁸, D-Arg²⁹]hGH-RH(1-29)NH₂

20 Peptide 22

[PhAc⁰, D-Arg², Phe(pCl)⁶, D-Abu⁸, Tyr(Me)¹⁰, Abu¹⁵, D-Arg²⁷, Arg²⁸, D-Arg²⁹]hGH-RH(1-29)NH₂ Peptide 23

[PhAc⁰, D-Arg², Phe(pCl)⁶, Lys⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 24

25 [PhAc⁰, D-Arg², Phe(pCl)⁶, Orn⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 25

[PhAc⁰, D-Arg², Phe(pCl)⁶, D-Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 26

[PhAc⁰, D-Arg², Phe(pCl)⁶, D-Har⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-

30 29)NH₂ Peptide 27

[PhAc⁰, D-Arg², Phe(pCl)⁶, D-Lys⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-

29)NH₂ Peptide 28

[PhAc⁰, D-Arg², Phe(pCl)⁶, D-Orn⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 29

[PhAc⁰, D-Arg², Phe(pCl)⁶, Cit⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 30

EXAMPLE III

Biological Activity

The peptides of the present invention were tested in in vitro and in vivo assays for their ability to inhibit the hGH-RH(1-29)NH₂ induced GH release.

Superfused Rat Pituitary System.

The analogs were tested in vitro in a test described earlier (S. Vigh and A.V. Schally, Peptides 5:241-347, 1984) with modification (Z. Rekasi and A.V. Schally, P.N.A.S. 90:2146-2149, 1993).

Briefly, the cells are preincubated with peptides for 9 minutes (3mL) at various concentrations. Immediately after the incubation, 1 nM hGH-RH(1-29)NH₂ is administered for 3 minutes (1mL) [0 minute response]. To check the duration of the antagonistic effect of the analogue, 1 nM hGH-RH(1-29)NH₂ is applied 30, 60, 90, and 120 minutes later for 3 minutes [30, 60, 90, 120 min responses]. Net integral values of the GH responses are evaluated. GH responses are compared to and expressed as percent of the original GH response induced by 1 nM GH-RH(1-29)NH₂. The effect of the new antagonists are compared to that of [Ac-Tyr¹,D-Arg²]hGH-RH(1-29)NH₂, the "Standard antagonist".

Growth Hormone Radio-immunoassay.

Rat GH levels in aliquots of undiluted and diluted superfusion samples were measured by double-antibody radioimmunoassay using materials

supplied by the National Hormone and Pituitary Program, Baltimore, Maryland. The results of RIA were analyzed with a computer program developed in our institute (V. Csernus and A.V. Schally, in *Neuroendocrine Research Methods*, Harwood Academic (Greenstein, B.D. ed., London, pp. 71-109, 1991), hereby incorporated by reference. Inter-assay variation was less than 15% and intra-assay variation was less than 10%.

GH-RH Binding Assay. A sensitive radioreceptor binding assay was developed to determine the binding characteristics of the antagonists of GH-RH (G. Halmos, A.V. Schally et al., Receptor **3**, 87-97, 1993), hereby incorporated by reference. The assay is based on binding of labelled $[\text{His}^1, \text{Nle}^{27}] \text{hGH-RH}(1-32)\text{NH}_2$ to rat anterior pituitary membrane homogenates. Iodinated derivatives of $[\text{His}^1, \text{Nle}^{27}] \text{hGH-RH}(1-32)\text{NH}_2$ are prepared by the chloramine-T method (F.C. Greenwood et al., Biochemistry **89**:114-123, 1963), hereby incorporated by reference. Pituitaries from male Sprague-Dawley rats (250-300 g) are used to prepare crude membranes. For saturation binding analyses, membrane homogenates are incubated with at least 6 concentrations of $[\text{His}^1, ^{125}\text{I-Tyr}^{10}, \text{Nle}^{27}] \text{hGH-RH}(1-32)\text{NH}_2$, ranging from 0.005 to 0.35 nM in the presence or absence of excess unlabelled peptide (1 μM).

The pellet is counted for radioactivity in a γ -counter. The affinities of the antagonist peptides tested to rat pituitary GH-RH receptors are determined in competitive binding experiments. The final binding affinities are estimated by K_i (dissociation constant of the inhibitor-receptor complex) and are determined by the Ligand PC and McPherson computer programs of Munson and Rodbard (P.J. Munson and D. Rodbard, Anal. Biochem. **107**, 220-239, 1980). Relative affinities compared to $[\text{Ac-Tyr}^1, \text{D-Arg}^2] \text{hGH-RH}(1-29)\text{NH}_2$, the Standard antagonist, are calculated as the ratio of K_i of the tested GH-RH antagonist to the K_i of the Standard antagonist.

In Vivo Tests.

GH-RH antagonistic effect of the analogs was tested on young male Sprague-Dawley rats (200-250 g). In each experiment, 5 groups of 7 animals each were used. The compounds (80 $\mu\text{g/kg}$) and GH-RH(1-29) NH_2 (3 $\mu\text{g/kg}$) were dissolved in 5.5% mannitol and given intravenously to the jugular vein of the rats under Nembutal anesthesia. The time elapsed between the administrations of the antagonist and GH-RH varied between groups, according to the following schedule. The first group of animals received GH-RH injection 5 min after the administration of the antagonist; for the second, third and fourth group of animals, the time interval elapsed between the injection of the antagonist and that of GH-RH was 15, 30, and 60 min, respectively. The control group was first injected with the solvent alone instead of the antagonist, followed by a GH-RH injection 5 min later.

0.4 mL blood samples were taken for GH RIA before the administration of the antagonist ("blood0"), and 5 min after the injection of GH-RH ("blood1"). The GH response in each group was calculated as $\text{rGH} = (\text{GH}_{\text{blood1}} / \text{GH}_{\text{blood0}})$, mean \pm S.E.M. of the individual differences. Relative inhibition of GH response (%) in each group treated was calculated as $100 \times (\text{rGH}_{\text{treated}} - 1) / (\text{rGH}_{\text{control}} - 1)$.

Results in vitro.

The results of the in vitro antagonistic activities tested in superfused rat pituitary system and binding assay are summarized in Table II and Table III, respectively. As it can be seen from these data, the substitutions present in the molecules cause an immense increase in receptor binding as well as in inhibition of GH release *in vitro* as compared to the standard antagonist. The most potent antagonist *in vitro*, Peptide 1, caused a complete inhibition of the GH-RH induced GH release for 90 min, under the standard test conditions. The first sign of the recovery of the GH-RH responsiveness was detected 120 min after the exposure to this analog.

TABLE II
Inhibition of GH Release in Superfused Rat Pituitary System

Antagonist	Dose (nM)	Inhibition of GH release (%)				
		0 min	30 min	60 min	90 min	120 min
Standard antagonist:	100	62.1	2.5	19		
Peptide 1	30	100	100	100	100	94
Peptide 2	30	100	100	100	100	91
Peptide 3	30	85	98	91	92	87
Peptide 4	30	83	86	80	79	68
Peptide 5	30	79	77	59	58	50
Peptide 6	30	93	93	97	95	90
Peptide 7	30	97	92	82	77	65
Peptide 8	30	100	100	98	97	90
Peptide 9	30	91	87	79	72	59
Peptide 10	30	75	69	46	24	22
Peptide 11	30	96	89	80	56	42
Peptide 12	30	68	75	48	52	52
Peptide 16	30	65	87	83	56	65
Peptide 19	30	58	47	59	23	39

TABLE III
K_i Values and Relative Affinities (R.A.) of hGH-RH Antagonists

Peptide	K _i (nM)	R.A.
Standard	2.94	1
Peptide 1	0.044	67
Peptide 2	0.046	64
Peptide 3	0.068	43
Peptide 4	0.087	34
Peptide 5	0.036	82
Peptide 9	0.058	51
Peptide 10	0.107	27
Peptide 11	0.071	41
Peptide 12	0.070	42
Peptide 16	0.070	42

Results in vivo.

Table IV shows the serum GH responses and their relative inhibitions in rats pretreated with GH-RH antagonists. All of the tested analogs (Peptide 1, Peptide 2, Peptide 3, Peptide 4, Peptide 8, Peptide 9, Peptide 11, and Peptide 16) produce strong and long-lasting inhibition of the GH release stimulated by hGH-RH(1-29)NH₂. Peptide 1 and Peptide 2 are the most

potent on the short term, inhibiting GH response by 95% and 91%, when given 5 min before the hGH-RH(1-29)NH₂. The effect of these two peptides lasts for at least 30 min. On the other hand, peptides Peptide 11 and Peptide 3, which are slightly less potent on the short term, are extremely long-acting: their effect persists for at least 60 min.

TABLE IV

Serum GH Responses and Relative Inhibitions of GH Responses in Rats Pretreated with GH-RH Antagonists at Different Time Intervals Prior to hGH-RH(1-29)NH₂ Administration

Antagonist	Time interval (min)	GH response mean \pm S.E.M.	Relative inhibition %
15	Peptide 1		
	5	1.13 \pm 0.13	95
	15	1.87 \pm 0.13	68
	30	1.97 \pm 0.16	64
	60	3.89 \pm 0.65	-8
20	control	3.68 \pm 0.78	0
25	Peptide 2		
	5	1.40 \pm 0.29	91
	15	3.45 \pm 0.13	45
	30	3.64 \pm 0.71	41
	60	5.41 \pm 1.35	2
30	control	5.47 \pm 0.97	0
35	Peptide 3		
	5	3.01 \pm 0.41	68
	15	3.63 \pm 0.80	61
	30	4.89 \pm 0.95	41
	60	5.36 \pm 0.63	34
40	control	7.65 \pm 0.66	0
45	Peptide 4		
	5	3.08 \pm 0.58	82
	15	7.73 \pm 1.12	43
	30	12.00 \pm 2.54	7
	60	11.88 \pm 0.90	8
50	control	12.82 \pm 1.38	0
55	Peptide 8		
	5	2.3 \pm 0.3	74
	15	3.6 \pm 0.4	46
	30	5.5 \pm 0.6	8
	60	5.8 \pm 0.7	1
60	control	5.8 \pm 0.5	0

5	Peptide 9	5	2.75 ± 0.75	75
		15	3.85 ± 0.50	59
		30	3.34 ± 0.30	67
		60	7.32 ± 1.16	10
10	Peptide 11	5	5.15 ± 1.11	84
		15	13.64 ± 1.41	50
		30	16.04 ± 4.36	40
		60	16.61 ± 6.02	38
		control	26.17 ± 3.92	0
15	Peptide 16	5	2.89 ± 0.65	77
		15	3.44 ± 0.62	70
		30	5.24 ± 1.36	48
		60	9.19 ± 1.89	0
		control	9.13 ± 1.69	0
20				

EXAMPLE IV

Oncological tests.

Antitumor activities of the peptides of the present invention were tested in various cancer models. The antitumor effects of these new peptides were compared with those of earlier analogs (MZ-4-71 and MZ-5-156, subject to US Patent 5,550,212 and US Patent application 08/642,472.

30 Effect of GH-RH antagonists on MXT mouse mammary tumors

Estrogen independent MXT tumors were transplanted sc. to female BDF mice. One day after transplantation, the mice were divided into groups of 10 animals each, and the treatment was started. The mice in groups 1, 2, 3, and 4 received single injections daily of various GH-RH antagonists sc. at 20 μ g dose per day for 18 days. In groups 5 and 6, the peptides were administered by Alzet osmotic pumps releasing a daily amount of 20 μ g peptide. Tumors were measured regularly, and tumor volume was calculated. The mice were sacrificed on day 18 and tumor weights were measured.

Results

Peptides Peptide 1 and Peptide 3 had similar strong inhibitory effect on MXT mouse mammary cancers. Treatment with MZ-5-156 also resulted in a significant inhibition of tumor growth, but its effect was weaker than that of Peptide 1 or Peptide 3 (see Table V and Figure I).

TABLE V
Effect of Treatment with GH-RH Antagonists on MXT Mouse Mammary Tumors

Group	Tumor volume (mm ³)		Tumor weights (mg)	Number of surviving mice
	On day 14	On day 18		
1. Control	4051 ± 1007	7040 ± 646	7269 ± 292	5
2. MZ-5-156	2717 ± 773	5368 ± 408*	4885 ± 480*	4
3. Peptide 1 daily inj.	2924 ± 654	4373 ± 381**	6964 ± 676	6
4. Peptide 3 daily inj.	1902 ± 349**	3465 ± 607**	5266 ± 906	8
5. Peptide 1 pump	1329 ± 327**	3403 ± 584**	4810 ± 645*	7
6. Peptide 3 pump	1688 ± 220**	4272 ± 295**	5939 ± 453	8

*p < 0.05

**p < 0.01

Effect of GH-RH antagonists on MDA-MB-468 human breast cancer xenografts in nude mice

Nude mice bearing MDA-MB-468 hormone-independent human breast cancer xenografts were divided into groups of 10 animals each. The treated groups received single daily s.c. injections of 20 µg of GH-RH antagonists. One group was treated with Peptide 1, a second group was treated with MZ-5-156 for comparison. The control group was injected with the vehicle solvent. The treatment was continued for 5 weeks. Tumors were measured once a week and tumor volume was calculated. The mice were sacrificed at the end of experiment and tumor weights were measured.

Results

Both peptides exerted significant tumor-inhibitory effects on MDA-MB-468 xenografts. In the group injected with Peptide 1, 4 tumors showed constant regression during the experiment. Likewise, MZ-5-156 caused

regression of 3 tumors. After 5 weeks of treatment these cancers regressed to small scar-like tissue remnants. Histological examination of these tissues revealed undifferentiated epithelial tumors with extensive necrosis and only a narrow marginal line of living tumor tissue. In contrast, all tumors in the 5 control animals progressed steadily. Final tumor volume and weight in the treated groups were significantly reduced (see Table VI and Figure II), Peptide 1 having a stronger effect.

TABLE VI

10 Effect of Treatment with GH-RH Antagonists on MDA-MB-468 Human Breast Cancer Xenografts in Nude Mice

Group	Final tumor volume (mm ³)	Tumor weight (mg)	Number of regressed tumors
15 Control	477.5 ± 41.2	440.7 ± 37.7	0
Peptide 1	82.4 ± 29.1**	64.0 ± 28.7**	4
MZ-5-156	104.4 ± 32.2**	77.7 ± 31.7**	3

20 *p < 0.05 **p < 0.01

Effect of GH-RH antagonists on HT-29 human colon cancer xenografts in nude mice

- 25 HT-29 human colon cancers were transplanted sc. into male nude mice. 19 days after transplantation, the mice were divided into groups of 10 animals each, and the treatment was started. The mice received single daily injections of various GH-RH antagonists sc. at 20 µg dose per day for 6 weeks. Tumors were measured regularly, and tumor volume was calculated.
- 30 The mice were sacrificed at the end of experiment and tumor weights were measured.

Results

- Peptides Peptide 1 and MZ-5-156 had equally strong inhibitory effect
- 35 on HT-29 human colon cancers. Treatment with Peptide 9 resulted in a smaller but still significant inhibition of tumor growth. Peptide 11 and MZ-4-71 had only little nonsignificant effect. (The results are summarized in Table

VII and Figure III).

TABLE VII
Effect of Treatment with GH-RH Antagonists on HT-29 Human Colon Cancer
5 Xenografts in Nude Mice

Group	Final tumor volume (mm ³)	Tumor weight (mg)
10 Control	2117 ± 751	2364 ± 835
MZ-4-71	1953 ± 400	2189 ± 458
MZ-5-156	908 ± 195*	1012 ± 174*
Peptide 11	1663 ± 610	1849 ± 681
Peptide 9	1194 ± 506*	1383 ± 576*
Peptide 1	890 ± 322*	1354 ± 480*
15		
*p < 0.05		

Effect of GH-RH antagonist Peptide 1 on U87MG human glioblastoma
20 xenografts in nude mice

Mice were implanted s.c. with U87MG glioblastomas and when tumors reached a volume of approx. 70 mm³ the mice were randomly divided into 2 experimental groups. One group was treated with Peptide 1 in single daily s.c. injections of 20 µg for 28 days, while the other group
25 served as control.

Results

Treatment with Peptide 1 inhibited the tumor growth by 77% versus the control group, after 4 weeks of treatment (see Table VIII and Figure IV).
30

TABLE VIII
Effect of Treatment with GH-RH Antagonist Peptide 1 on U87MG Human Glioblastoma Xenografts in Nude Mice

Group	Final tumor volume (mm ³)	Tumor weight (g)
35 Control	3425 ± 723	4.7 ± 1.3
Peptide 1	817 ± 323**	1.4 ± 0.7**

40 **p < 0.01

Effect of GH-RH antagonists on PC-3 human prostate cancer xenografts in nude mice

Male nude mice were implanted s.c. with 3 mm³ pieces of PC-3 human hormone-independent prostate cancer tissue into both flanks. When tumors reached a volume of approx. 40-50 mm³ the mice were divided into 3 experimental groups. The first and second groups were treated with peptides Peptide 3 and MZ-5-156, respectively, in single daily s.c. injections of 20 µg for 21 days, while the third group served as control. Tumor volumes were measured at weekly intervals, tumor weights were measured at the end of experiment.

Results

Both GH-RH antagonists inhibited growth of PC-3 tumors (see Table IX and Figure V). Peptide 3 exerted stronger growth inhibition (65% inhibition in tumor volume and 62% in tumor weight) than MZ-5-156 (52% and 46%, respectively).

TABLE IX

Effect of Treatment with GH-RH Antagonists on PC-3 Human Prostate Cancer Xenografts in Nude Mice

Group	Final tumor volume (mm ³)	Tumor weight (mg)
Control	307.9 ± 64.5	191.8 ± 42.6
Peptide 3	107.9 ± 12.5*	73.7 ± 11.9*
MZ-5-156	148.9 ± 41	104.3 ± 27.2

*p < 0.05

CLAIMS:

1. A peptide selected from the group having the formulae:
 $X-R^1-R^2-Asp-Ala-R^5-R^6-Thr-R^8-R^9-R^{10}-Arg-R^{12}-R^{13}-R^{14}-R^{15}-R^{16}-Leu-R^{18}-R^{19}-Arg-R^{21}-R^{22}-Leu-Gln-Asp-Ile-R^{27}-R^{28}-R^{29}-NH_2$
- 5 wherein X is PhAc, IndAc, Ibu, Nac, 1- or 2-Npr, or Fpr,
 R^1 is Tyr or His,
 R^2 is D-Arg or D-Cit,
 R^5 is Ile or Val,
 R^6 is Phe, Nal or Phe(Y), in which Y = F, Cl, Br,
10 R^8 is Asn, Gln, Ser, Thr, Ala, D-Asn, D-Gln, D-Ser, D-Thr, Abu, D-
 Abu, or Aib,
 R^9 is Arg, Har, Lys, Orn, D-Arg, D-Har, D-Lys, D-Orn, Cit, Nle, Tyr
 (Me), Ser, Ala or Aib,
 R^{10} is Tyr or Phe(Y), in which Y = H, F, Cl, Br, or OCH_3 ,
15 R^{12} is Lys, D-Lys, or Orn,
 R^{13} is Val or Nle,
 R^{14} is Leu or Nle,
 R^{15} is Gly, Ala, Abu, Nle or Gln,
 R^{16} is Gln or Arg,
20 R^{18} is Ser or Nle,
 R^{19} is Ala or Abu,
 R^{21} is Lys or Orn,
 R^{22} is Leu, Ala or Aib,
 R^{27} is Met, Leu, Nle, Abu, or D-Arg,
25 R^{28} is Arg, D-Arg, Ser, Asn, Asp, Ala or Abu,
 R^{29} is Arg, D-Arg, Har or D-Har,
 and pharmaceutically acceptable salts thereof.
2. A compound of claim 1 selected from the group consisting of
30 {PhAc⁹, D-Arg², Phe(pCl)⁶, Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹}hGH-RH(1-29)
NH₂ Peptide 1

[IndAc⁰, D-Arg², Phe(pCl)⁶, Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 2

[PhAc⁰, D-Arg², Phe(pCl)⁶, Har⁹, Tyr(Me)¹⁰, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 3

5 [PhAc⁰, D-Arg², Phe(pCl)⁶, Arg⁹, Tyr(Me)¹⁰, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 6

[PhAc⁰, His¹, D-Arg², Phe(pCl)⁶, Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 7

10 [Nac⁰, His¹, D-Arg², Phe(pCl)⁶, Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 8

3. A compound of claim 1 having the formula [PhAc⁰, D-Arg², Phe(pCl)⁶, Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 1

15 4. A compound of claim 1 having the formula [IndAc⁰, D-Arg², Phe(pCl)⁶, Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 2

5. A compound of claim 1 having the formula [PhAc⁰, D-Arg², Phe(pCl)⁶, Har⁹, Tyr(Me)¹⁰, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂
20 Peptide 3

6. A compound of claim 1 having the formula [PhAc⁰, D-Arg², Phe(pCl)⁶, Arg⁹, Tyr(Me)¹⁰, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂
25 Peptide 6

7. A compound of claim 1 having the formula [PhAc⁰, His¹, D-Arg², Phe(pCl)⁶, Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂
Peptide 7

30 8. A compound of claim 1 having the formula [Nac⁰, His¹, D-Arg², Phe(pCl)⁶, Arg⁹, Abu¹⁵, Nle²⁷, D-Arg²⁸, Har²⁹]hGH-RH(1-29)NH₂ Peptide 8

9. A method of suppressing excessive levels of GH in a patient in need of same which comprises administering said patient an effective amount of a compound of claim 1.

5 10. A method of treating a patient having a cancer carrying receptors for IGF-I or -II which comprises administering to said patient an effective amount of a compound of claim 1.

11. A method for inhibiting IGF-II levels in tumors (cancers) and the
10 expression of mRNA for IGF-II in the same tumors, which comprises administering to said patient an effective amount of a compound of claim 1.

ABSTRACT

There is provided a novel series of synthetic analogs of hGH-RH(1-29)NH₂. These analogs inhibit the activity of endogenous hGH-RH, and therefore prevent the release of growth hormone. The stronger inhibitory potencies of the new analogs, as compared to previously described ones, results from replacement of various amino acids.

10

d:\c_drive\wp51\shal\30b5.spc

FIGURE I

Antitumor Effects of GH-RH Antagonists Peptide 1, Peptide 3, and MZ-5-156 on MXT Mouse Mammary Cancers




FIGURE II
Antitumor Effects of GH-RH Antagonists Peptide 1, and MZ-5-156 on
MDA-MB-468 Human Breast Cancer Xenografts in Nude Mice




FIGURE III

Antitumor Effects of GH-RH Antagonists Peptide 1, Peptide 9, Peptide 11, MZ-4-71, and MZ-5-156 on HT-29 Human Colon Cancer Xenografts in Nude Mice




FIGURE IV
Antitumor Effect of GH-RH Antagonist Peptide 1 on U87MG Human Glioblastoma Xenografts in Nude Mice





FIGURE V
Antitumor Effects of GH-RH Antagonists Peptide 3, and MZ-5-156 on
PC-3 Human Prostate Cancer Xenografts in Nude Mice



* $p = 0.056$

** $p = 0.013$

Declaration and Power of Attorney For Patent Application English Language Declaration

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled
ANTAGONISTIC ANALOGS OF GB-RH INHIBITING IGF-I AND-II

the specification of which

(check one)

☒ is attached hereto.

☐ was filed on _____ as

Application Serial No. _____

and was amended on _____
(if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, §1.56(a).

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Prior Foreign Application(s)

Priority Claimed

(Number) _____ (Country) _____ (Day/Month/Year Filed) _____

☐ Yes ☐ No

(Number) _____ (Country) _____ (Day/Month/Year Filed) _____

☐ Yes ☐ No

(Number) _____ (Country) _____ (Day/Month/Year Filed) _____

☐ Yes ☐ No

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

(Application Serial No.)	(Filing Date)	(Status) (patented, pending, abandoned)
(Application Serial No.)	(Filing Date)	(Status) (patented, pending, abandoned)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. (*list name and registration number*)

Omri M. Behr, Esq.; Regis. No. 22,940

Send Correspondence to:

Omri M. Behr, Esq., Behr 7 Adams, 325 Pierson Avenue, Edison, NJ 08837

Direct Telephone Calls to: (*name and telephone number*)

Omri M. Behr, Esq., 1-732-494-5240

Full name of sole or first inventor ANDREW V. SCHALLY	
Inventor's signature <i>Andrew V. Schally</i>	Date NOVEMBER 16, 1998
Residence Metairie, LA 70006	
Citizenship United States	
Post Office Address 5025 Kawanee Avenue, Metairie, LA 70006	
Full name of second joint inventor, if any JOZSEF VARGA	
Second inventor's signature <i>Josef L. Varga</i>	Date NOVEMBER 16, 1998
Residence New Orleans, LA 70125	
Citizenship Hungary	
Post Office Address 6440 So. Claiborne Avenue - Apt. 203, New Orleans, LA 70125	

(Supply similar information and signature for third and subsequent joint inventors.)

Full name of <input type="checkbox"/> Inventor	
MARTA ZARANDI	
Inventor's signature <i>Marta Zarandi</i>	
Date 16/11/98	
Residence Szeged H. 6723, Hungary	
Citizenship Hungary	
Post Office Address Dugonics H. 25., Szeged H. 6724, Hungary	
Full name of second joint inventor, if any	
Second inventor's signature	
Date	
Residence	
Citizenship	
Post Office Address	

Full name of <input type="checkbox"/> Inventor	
Inventor's signature	
Date	
Residence	
Citizenship	
Post Office Address	

Full name of <input type="checkbox"/> joint inventor, if any	
Second inventor's signature	
Date	
Residence	
Citizenship	
Post Office Address	

(Supply similar information and signature for third and subsequent joint inventors.)